

Domination Value in $P_2 \square P_n$ and $P_2 \square C_n$

Eunjeong Yi

Texas A&M University at Galveston
Galveston, TX 77553, USA
yie@tamug.edu

March 2, 2012

Abstract

A set $D \subseteq V(G)$ is a *dominating set* of a graph G if every vertex of G not in D is adjacent to at least one vertex in D . A *minimum dominating set* of G , also called a $\gamma(G)$ -set, is a dominating set of G of minimum cardinality. For each vertex $v \in V(G)$, we define the *domination value* of v to be the number of $\gamma(G)$ -sets to which v belongs. In this paper, we find the total number of minimum dominating sets and characterize the domination values for $P_2 \square P_n$ and $P_2 \square C_n$.

1 Introduction

Let $G = (V(G), E(G))$ be a simple, undirected, and nontrivial graph. For $S \subseteq V(G)$, we denote by $\langle S \rangle$ the subgraph of G induced by S . For a vertex $v \in V(G)$, the *open neighborhood* of v is the set $N(v) = \{u \mid uv \in E(G)\}$, and the *closed neighborhood* of v is the set $N[v] = N(v) \cup \{v\}$. For $S \subseteq V(G)$, the *open neighborhood* of S is the set $N(S) = \cup_{v \in S} N(v)$ and the *closed neighborhood* of S is the set $N[S] = N(S) \cup S$.

A set $D \subseteq V(G)$ is a *dominating set* if $N[D] = V(G)$, and is a *total dominating set* if $N(D) = V(G)$. The *domination number* of a graph G , denoted by $\gamma(G)$, is the minimum of the cardinalities of all dominating sets of G . A *minimum dominating set* of G , also called a $\gamma(G)$ -set, is a dominating set of G of minimum cardinality. For discussions on domination (resp. total domination) in graphs, see [1, 2, 6, 9, 10, 17] (resp. see [5, 9, 12]). Slater [18] introduced the notion of the number of dominating sets of G , which he denoted by $\text{HED}(G)$ in honor of Steve Hedetniemi on

the occasion of his 60th birthday; further, Slater used $\#\gamma(G)$ to denote the number of $\gamma(G)$ -sets. Following [14, 19], we denote by $\tau(G)$ the total number of $\gamma(G)$ -sets. For each vertex $v \in V(G)$, we define the *domination value* of v in G , denoted by $DV_G(v)$, to be the number of $\gamma(G)$ -sets to which v belongs; we often drop G when ambiguity is not a concern. Clearly, $0 \leq DV_G(v) \leq \tau(G)$ for any graph G and for any vertex $v \in V(G)$. See [19] for an introductory discussion on domination value in graphs and [14] for an introductory discussion on total domination value in graphs.

The *Cartesian product* of two graphs G and H , denoted by $G \square H$, is the graph with the vertex set $V(G) \times V(H)$ such that (u, v) is adjacent to (u', v') if and only if (i) $u = u'$ and $vv' \in E(H)$ or (ii) $v = v'$ and $uu' \in E(G)$. For other graph theory terminology, refer to [4].

We denote by P_n and C_n the path and the cycle on n vertices, respectively. In [13], Jacobson and Kinch obtained the results on $\gamma(P_m \square P_n)$ for $m = 2, 3, 4$. Later, Hare developed an algorithm to compute $\gamma(P_m \square P_n)$ and was able to find expressions for $\gamma(P_m \square P_n)$ for a number of different values of m and n (see [8]). Chang and Clark proved the formulas found by Hare for $\gamma(P_5 \square P_n)$ and $\gamma(P_6 \square P_n)$ in [3]. The complexity of determining $\gamma(P_m \square P_n)$ is open as of [11]. In [15], Klavžar and Seifter obtained results on $\gamma(C_m \square C_n)$ for $m = 3, 4, 5$.

In section 2, we present relevant results from [19]. In sections 3 and 4, noting $\gamma(P_2 \square P_n) \neq \gamma(P_2 \square C_n)$ for $n \equiv 0 \pmod{4}$, we investigate the total number of minimum dominating sets and the domination value for two classes of graphs, $P_2 \square P_n$ and $P_2 \square C_n$.

2 Preliminaries and domination value in paths and cycles

We first recall the following observations.

Observation 2.1. [19] $\sum_{v \in V(G)} DV_G(v) = \tau(G) \cdot \gamma(G)$

Observation 2.2. [19] If there is an isomorphism of graphs carrying a vertex v in G to a vertex v' in G' , then $DV_G(v) = DV_{G'}(v')$.

It is well known that $\gamma(P_n) = \gamma(C_n) = \lceil \frac{n}{3} \rceil$. If we let the vertices of the path P_n be labeled 1 through n consecutively, then we have the following

Theorem 2.3. [19] For $n \geq 2$,

$$\tau(P_n) = \begin{cases} 1 & \text{if } n \equiv 0 \pmod{3} \\ n + \frac{1}{2} \lfloor \frac{n}{3} \rfloor (\lfloor \frac{n}{3} \rfloor - 1) & \text{if } n \equiv 1 \pmod{3} \\ 2 + \lfloor \frac{n}{3} \rfloor & \text{if } n \equiv 2 \pmod{3}. \end{cases}$$

For the domination value of a vertex v on P_n , by Observation 2.2, $DV(v) = DV(n+1-v)$ for $1 \leq v \leq n$. More precisely, we have the classification results which follow.

Corollary 2.4. [19] Let $v \in V(P_{3k})$, where $k \geq 1$. Then

$$DV(v) = \begin{cases} 0 & \text{if } v \equiv 0, 1 \pmod{3} \\ 1 & \text{if } v \equiv 2 \pmod{3}. \end{cases}$$

Proposition 2.5. [19] Let $v \in V(P_{3k+1})$, where $k \geq 1$. Write $v = 3q + r$, where q and r are non-negative integers such that $0 \leq r < 3$. Then, noting $\tau(P_{3k+1}) = \frac{1}{2}(k^2 + 5k + 2)$, we have

$$DV(v) = \begin{cases} \frac{1}{2}q(q+3) & \text{if } v \equiv 0 \pmod{3} \\ (q+1)(k-q+1) & \text{if } v \equiv 1 \pmod{3} \\ \frac{1}{2}(k-q)(k-q+3) & \text{if } v \equiv 2 \pmod{3}. \end{cases}$$

Proposition 2.6. [19] Let $v \in V(P_{3k+2})$, where $k \geq 0$. Write $v = 3q + r$, where q and r are non-negative integers such that $0 \leq r < 3$. Then, noting $\tau(P_{3k+2}) = k + 2$, we have

$$DV(v) = \begin{cases} 0 & \text{if } v \equiv 0 \pmod{3} \\ 1 + q & \text{if } v \equiv 1 \pmod{3} \\ k + 1 - q & \text{if } v \equiv 2 \pmod{3}. \end{cases}$$

If we let the vertices of the cycle C_n be labeled 1 through n cyclically, then we have the following

Theorem 2.7. [19] For $n \geq 3$,

$$\tau(C_n) = \begin{cases} 3 & \text{if } n \equiv 0 \pmod{3} \\ n(1 + \frac{1}{2} \lfloor \frac{n}{3} \rfloor) & \text{if } n \equiv 1 \pmod{3} \\ n & \text{if } n \equiv 2 \pmod{3}. \end{cases}$$

By Theorem 2.7, Observation 2.1, Observation 2.2, and the vertex-transitivity of C_n , we have the following

Corollary 2.8. [19] Let $v \in V(C_n)$, where $n \geq 3$. Then

$$DV(v) = \begin{cases} 1 & \text{if } n \equiv 0 \pmod{3} \\ \frac{1}{2} \lceil \frac{n}{3} \rceil (1 + \lceil \frac{n}{3} \rceil) & \text{if } n \equiv 1 \pmod{3} \\ \lceil \frac{n}{3} \rceil & \text{if } n \equiv 2 \pmod{3}. \end{cases}$$

3 Total number of minimum dominating sets and domination value in $P_2 \square P_n$

We consider $P_2 \square P_n$ ($n \geq 2$) as two copies of P_n with vertices labeled x_1, x_2, \dots, x_n and y_1, y_2, \dots, y_n with only the edges $x_i y_i$, for each i ($1 \leq i \leq n$), between two paths (see Figure 1).




Figure 1: Labeling of vertices of $P_2 \square P_n$

We first recall the following.

Theorem 3.1. [13] For $n \geq 2$, $\gamma(P_2 \square P_n) = \lceil \frac{n+1}{2} \rceil$.

Lemma 3.2. Let $G = P_2 \square P_n$, where $n \geq 2$. If neither x_1 nor y_1 belongs to a $\gamma(G)$ -set D , then $\{x_2, y_2\} \subseteq D$. (Likewise, if neither x_n nor y_n belongs to D , then $\{x_{n-1}, y_{n-1}\} \subseteq D$.)

Proof. By definition of a dominating set, either x_1 or a vertex in $N(x_1) = \{x_2, y_1\}$ belongs to D . If $x_1 \notin D$ and $y_1 \notin D$, then $x_2 \in D$. Similarly, either $y_1 \in D$ or a vertex in $N(y_1) = \{x_1, y_2\}$ belongs to D . If $x_1 \notin D$ and $y_1 \notin D$, then $y_2 \in D$ as well. Thus $x_1 \notin D$ and $y_1 \notin D$ implies $\{x_2, y_2\} \subseteq D$. \square

Lemma 3.3. Let $G = P_2 \square P_n$, where $n \geq 3$. If there exists a $\gamma(G)$ -set containing no vertex of degree two, then $n = 3$ or $n = 6$.

Proof. Suppose that D is a $\gamma(G)$ -set such that $\{x_1, y_1, x_n, y_n\} \cap D = \emptyset$. Let $S_0 = \{x_2, y_2, x_{n-1}, y_{n-1}\}$. Then, by Lemma 3.2, $S_0 \subseteq D$. Note that $|S_0| = 2$ if and only if $n = 3$: in this case, $\gamma(P_2 \square P_3) = 2$ and $S_0 = \{x_2, y_2\}$ is a $\gamma(P_2 \square P_3)$ -set. If $4 \leq n \leq 5$, then $|S_0| = 4$ and $\gamma(P_2 \square P_n) = 3$, and thus $S_0 \not\subseteq D$. If $n = 6$, then $|S_0| = 4$ and $\gamma(P_2 \square P_6) = 4$: in fact, $S_0 = \{x_2, y_2, x_5, y_5\}$ is a $\gamma(P_2 \square P_6)$ -set. Now, we need to consider $n \geq 7$. Suppose that $S_0 \subseteq D$; we consider two cases.

Case 1. $n = 2k$, where $k \geq 4$: Here, $\gamma(P_2 \square P_{2k}) = k + 1$. Since $N[S_0] = \{x_i, y_i \mid 1 \leq i \leq 3\} \cup \{x_j, y_j \mid 2k-2 \leq j \leq 2k\}$, the part of $P_2 \square P_{2k}$ not dominated by S_0 is a $P_2 \square P_{2k-6}$. So, $k - 3$ vertices of $D - S_0$ must dominate $P_2 \square P_{2k-6}$. But $\gamma(P_2 \square P_{2k-6}) = k - 2$ by Theorem 3.1, and we reach a contradiction.

Case 2. $n = 2k + 1$, where $k \geq 3$: Here, $\gamma(P_2 \square P_{2k+1}) = k + 1$. Since $N[S_0] = \{x_i, y_i \mid 1 \leq i \leq 3\} \cup \{x_j, y_j \mid 2k - 1 \leq j \leq 2k + 1\}$, the part of $P_2 \square P_{2k+1}$ not dominated by S_0 is a $P_2 \square P_{2k-5}$. So, $k - 3$ vertices of $D - S_0$ must dominate $P_2 \square P_{2k-5}$. But $\gamma(P_2 \square P_{2k-5}) = k - 2$ by Theorem 3.1, and we reach a contradiction.

Thus, we have shown that if $S_0 \subseteq D$, then $n = 3$ or $n = 6$. \square

Next we compute the total number of $\gamma(P_2 \square P_n)$ -sets for $n \geq 2$.

Theorem 3.4. *For $n \geq 2$,*

$$\tau(P_2 \square P_n) = \begin{cases} 6 & \text{if } n = 2 \\ 3 & \text{if } n = 3 \\ 17 & \text{if } n = 6 \\ 2 & \text{if } n \text{ is odd and } n \neq 3 \\ 2n + 4 & \text{if } n \text{ is even and } n \neq 2, 6. \end{cases}$$

Proof. Let D be a $\gamma(P_2 \square P_n)$ -set for $n \geq 2$. Notice that no D contains both x_1 and y_1 , or both x_n and y_n , unless $n = 2$. We consider two cases.

Case 1. $n \geq 3$ is odd: Here, $\gamma(P_2 \square P_n) = \frac{n+1}{2}$. By Lemma 3.3, if there is a D containing no vertex of degree two then $n = 3$. Moreover, we note that $\{x_2, y_2\} \subseteq D$ if and only if $n = 3$: If $\{x_2, y_2\} \subseteq D$ and $n > 3$, then the part of $P_2 \square P_n$ not dominated by $\{x_2, y_2\}$ is a $P_2 \square P_{n-3}$, and $\frac{n-3}{2}$ vertices of $D - \{x_2, y_2\}$ must dominate $P_2 \square P_{n-3}$. But $\gamma(P_2 \square P_{n-3}) = \frac{n-1}{2}$ by Theorem 3.1, and we reach a contradiction. So, if $n > 3$, by Lemma 3.2, either $x_1 \in D$ or $y_1 \in D$. One can easily check that $x_1 \in D$ uniquely determines a γ -set $D = \{x_i, y_j \mid i \equiv 1, j \equiv 3 \pmod{4}\}$. Similarly, $y_1 \in D$ uniquely determines a γ -set $D = \{x_i, y_j \mid i \equiv 3, j \equiv 1 \pmod{4}\}$. Thus, $\tau(P_2 \square P_n) = 2$ for $n \neq 3$, and $\tau(P_2 \square P_3) = 3$ by Lemma 3.3. (See Figure 2 for the three $\gamma(P_2 \square P_3)$ -sets, where the solid black vertices in each $P_2 \square P_3$ form a $\gamma(P_2 \square P_3)$ -set.)




Figure 2: γ -sets for $P_2 \square P_3$

Case 2. $n \geq 2$ is even: Here, $\gamma(P_2 \square P_n) = \frac{n}{2} + 1$. If $n = 2$, then $\gamma(P_2 \square P_2) = 2$ and $\tau(P_2 \square P_2) = \tau(C_4) = \binom{4}{2} = 6$. We consider $n \geq 4$. By Lemma 3.3, if there is a D containing no vertex of degree two (i.e., $\{x_2, y_2, x_{n-1}, y_{n-1}\} \subseteq D$), then $n = 6$. We consider three subcases.

Subcase 2.1. $\{x_2, y_2\} \subseteq D$ and $\{x_{n-1}, y_{n-1}\} \cap D = \emptyset$: Let τ_1 be the number of such $\gamma(P_2 \square P_n)$ -sets for $n \geq 4$. Note that the part of $P_2 \square P_n$ not dominated by $\{x_2, y_2\}$ is a $P_2 \square P_{n-3}$. So, τ_1 equals the number of $\gamma(P_2 \square P_{n-3})$ -sets with $\gamma(P_2 \square P_{n-3}) = \frac{n}{2} - 1$. One can easily see that $\tau_1 = 2$ when $n = 4, 6$. Since $\tau_1(P_2 \square P_{n-3}) = 2$ for $n \geq 8$ by Case 1, we have $\tau_1 = 2$ for $n \geq 4$.

Subcase 2.2. $\{x_2, y_2\} \cap D = \emptyset$ and $\{x_{n-1}, y_{n-1}\} \subseteq D$: Let τ_2 be the number of such $\gamma(P_2 \square P_n)$ -sets for $n \geq 4$. By Observation 2.2 and Subcase 2.1, we have $\tau_2 = 2$ for $n \geq 4$.

Subcase 2.3. $\{x_2, y_2\} \not\subseteq D$ and $\{x_{n-1}, y_{n-1}\} \not\subseteq D$: By Lemma 3.2, $|\{x_1, y_1\} \cap D| = 1$ and $|\{x_n, y_n\} \cap D| = 1$. Let D (resp. D') be such a γ -set of $G = P_2 \square P_n$ (resp. $G' = P_2 \square P_{n+2}$), where $n \geq 4$. And let τ_3 (resp. τ'_3) be the number of such γ -sets of G (resp. G'). We will show that $\tau_3 = 2n$, for $n \geq 4$, using induction. The base case, $n = 4$, is easily verified (see Figure 3). Assume that $\tau_3 = 2n$ for $n \geq 4$. If $x_1 \in D$, then each D extends to D' such that $D' = D \cup \{x_{n+2}\}$ if $y_n \in D$ and $D' = D \cup \{y_{n+2}\}$ if $x_n \in D$; in addition, there are two additional $\gamma(G')$ -sets which do not come from any $\gamma(G)$ -sets, i.e., $\{x_i, y_j \mid i \equiv 1, j \equiv 3 \pmod{4} \text{ and } 1 \leq i, j \leq n+1\} \cup \{x_{n+2}\}$ and $\{x_i, y_j \mid i \equiv 1, j \equiv 3 \pmod{4} \text{ and } 1 \leq i, j \leq n+1\} \cup \{y_{n+2}\}$. Similarly, if $y_1 \in D$, then each D extends to D' and there are two additional $\gamma(G')$ -sets which do not come from $\gamma(G)$ -sets. So, $\tau'_3 = \tau_3 + 4 = 2n + 4 = 2(n + 2)$.

Now, noting that $\{x_2, y_2, x_{n-1}, y_{n-1}\} \subseteq D$ implies $n = 6$, combine the three disjoint cases to get $\tau = \tau_1 + \tau_2 + \tau_3 = 2 + 2 + 2n = 2n + 4$ if $n \neq 2, 6$ and $\tau(P_2 \square P_6) = (2 \cdot 6 + 4) + 1 = 17$. \square

See Figure 3 for the collection of $\gamma(P_2 \square P_4)$ -sets, where the solid black vertices in each $P_2 \square P_4$ form a $\gamma(P_2 \square P_4)$ -set.




Figure 3: γ -sets for $P_2 \square P_4$

As an immediate consequence of Theorem 3.4 for an odd $n \geq 3$, we have the following

Corollary 3.5. *Let $n \geq 3$ be an odd number.*

(i) For each $v \in V(P_2 \square P_3)$, $DV(v) = 1$.

(ii) For $x_i, y_i \in V(P_2 \square P_n)$, where $n \geq 5$,

$$DV(x_i) = DV(y_i) = \begin{cases} 1 & \text{if } i \text{ is odd} \\ 0 & \text{if } i \text{ is even} . \end{cases}$$

Proposition 3.6. Let $n \geq 2$ be an even number.

(i) For each $v \in V(P_2 \square P_2)$, $DV(v) = 3$.

(ii) For $x_i, y_i \in V(P_2 \square P_n)$, where $n \geq 4$ and $n \neq 6$,

$$DV(x_i) = DV(y_i) = \begin{cases} n+2-i & \text{if } i \text{ is odd and } 1 \leq i \leq n-3 \\ 4 & \text{if } i = 2 \text{ or } i = n-1 \\ i+1 & \text{if } i \text{ is even and } 4 \leq i \leq n . \end{cases} \quad (1)$$

(iii) For $x_i, y_i \in V(P_2 \square P_6)$,

$$DV(x_i) = DV(y_i) = \begin{cases} 7 & \text{if } i = 1 \text{ or } i = 6 \\ 5 & \text{if } 2 \leq i \leq 5 . \end{cases} \quad (2)$$

Proof. Let $n \geq 2$ be an even number.

(i) Note that $P_2 \square P_2 \cong C_4$, $\gamma(C_4) = 2$, and $\tau(C_4) = 6$. By Observation 2.1, Observation 2.2, and the vertex-transitivity, $DV(v) = 3$ for each $v \in V(P_2 \square P_2)$.

(ii) For an even $n \geq 4$, let D (resp. D') be a γ -set of $G = P_2 \square P_n$ (resp. $G' = P_2 \square P_{n+2}$). Since $DV_G(x_i) = DV_G(y_i)$ for each i ($1 \leq i \leq n$), it suffices to compute $DV_G(x_i)$ for $1 \leq i \leq n$. We consider two cases.

Case 1. $\{x_1, y_1\} \cap D = \emptyset$: By Lemma 3.2, $\{x_2, y_2\} \subseteq D$. Denote by $DV^1(v)$ the number of such D 's containing v . Notice that there are two such $\gamma(G)$ -sets. We will show, by induction, that

$$DV_G^1(x_i) = \begin{cases} 2 & \text{if } i = 2 \\ 1 & \text{if } i \geq 4 \text{ and } i \text{ is even} \\ 0 & \text{if } i \text{ is odd} . \end{cases} \quad (3)$$

For $n = 4$ (the base case), the two γ -sets are $\{x_2, y_2, x_4\}$ and $\{x_2, y_2, y_4\}$, thus satisfying (3). Assume that (3) holds for G . Let D_1 and D_2 be $\gamma(G)$ -sets, containing both x_2 and y_2 , such that $x_n \in D_1$ and $y_n \in D_2$. Then D_1 extends to $D'_1 = D_1 \cup \{y_{n+2}\}$ and D_2 extends to $D'_2 = D_2 \cup \{x_{n+2}\}$,

where D'_1 and D'_2 are $\gamma(G')$ -sets. So, $DV_{G'}^1(x_i) = DV_G^1(x_i)$ for $1 \leq i \leq n$, $DV_{G'}^1(x_{n+1}) = 0$, and $DV_{G'}^1(x_{n+2}) = 1$. Thus

$$DV_{G'}^1(x_i) = \begin{cases} 2 & \text{if } i = 2 \\ 1 & \text{if } i \geq 4 \text{ and } i \text{ is even} \\ 0 & \text{if } i \text{ is odd,} \end{cases}$$

proving (3).

Case 2. $x_1 \in D$ or $y_1 \in D$: Denote by $DV^2(v)$ the number of such D 's containing v . By Subcase 2.2 and Subcase 2.3 in the proof of Theorem 3.4, there are $2n + 2$ such $\gamma(G)$ -sets; $n + 1$ such D 's containing x_1 , and $n + 1$ such D 's containing y_1 . We will show, by induction, that

$$DV_G^2(x_i) = \begin{cases} i & \text{if } i \equiv 0, 2 \pmod{4} \text{ and } 2 \leq i \leq n \\ n + 2 - i & \text{if } i \equiv 1, 3 \pmod{4} \text{ and } 1 \leq i \leq n - 3 \\ 4 & \text{if } i = n - 1. \end{cases} \quad (4)$$

Noting that no $\gamma(G)$ -set contains both x_1 and y_1 , we consider two subcases.

Subcase 2.1. $x_1 \in D$: Denote by $DV^{2,1}(v)$ the number of such D 's containing v . For $n = 4$ (the base case), one can check that there are five such γ -sets: $\{x_1, x_2, y_4\}$, $\{x_1, y_2, x_4\}$, $\{x_1, y_3, x_4\}$, $\{x_1, y_3, y_4\}$, and $\{x_1, x_3, y_3\}$. Let D_1, D_2, \dots, D_{n+1} be $\gamma(G)$ -sets containing x_1 , where $\{x_{n-1}, y_{n-1}\} \subseteq D_{n+1}$. Then, for $1 \leq i \leq n$, each D_i extends to $D'_i = D_i \cup \{x_{n+2}\}$ if $y_n \in D_i$ and $D'_i = D_i \cup \{y_{n+2}\}$ if $x_n \in D_i$, where each D'_i ($1 \leq i \leq n$) is a $\gamma(G')$ -set; $D_{n+1} = \{x_i, y_j \mid i \equiv 1, j \equiv 3 \pmod{4} \text{ and } 1 \leq i, j \leq n - 2\} \cup \{x_{n-1}, y_{n-1}\}$ does not extend to a $\gamma(G')$ -set, but there exists a $\gamma(G')$ -set $D'_{n+1} = \{x_i, y_j \mid i \equiv 1, j \equiv 3 \pmod{4} \text{ and } 1 \leq i, j \leq n\} \cup \{x_{n+1}, y_{n+1}\}$ which does not come from any $\gamma(G)$ -set. Further, there exist two additional $\gamma(G')$ -sets which do not come from any $\gamma(G)$ -sets such as $D'_{n+2} = \{x_i, y_j \mid i \equiv 1, j \equiv 3 \pmod{4} \text{ and } 1 \leq i, j \leq n + 1\} \cup \{x_{n+2}\}$ and $D'_{n+3} = \{x_i, y_j \mid i \equiv 1, j \equiv 3 \pmod{4} \text{ and } 1 \leq i, j \leq n + 1\} \cup \{y_{n+2}\}$. So, noting that n is even, we have the following:

$$DV_{G'}^{2,1}(x_i) = \begin{cases} DV_G^{2,1}(x_i) & \text{if } i \equiv 0, 2, 3 \pmod{4} \text{ and } 1 \leq i \leq n - 2 \\ DV_G^{2,1}(x_i) + 2 & \text{if } i \equiv 1 \pmod{4} \text{ and } 1 \leq i \leq n - 2, \end{cases}$$

$$DV_{G'}^{2,1}(x_{n-1}) = \begin{cases} DV_G^{2,1}(x_{n-1}) - 1 & \text{if } n \equiv 0 \pmod{4} \\ DV_G^{2,1}(x_{n-1}) + 2 & \text{if } n \equiv 2 \pmod{4}, \end{cases}$$

$$DV_{G'}^{2,1}(x_{n+1}) = \begin{cases} 3 & \text{if } n \equiv 0 \pmod{4} \\ 1 & \text{if } n \equiv 2 \pmod{4}, \end{cases}$$

$$DV_{G'}^{2,1}(x_n) = DV_G^{2,1}(x_n), \text{ and } DV_{G'}^{2,1}(x_{n+2}) = \frac{n}{2} + 1.$$

Subcase 2.2. $y_1 \in D$: Denote by $DV^{2,2}(v)$ the number of such D 's containing v . For $n = 4$ (the base case), one can check that there are five such γ -sets: $\{y_1, y_2, x_4\}$, $\{y_1, x_2, y_4\}$, $\{y_1, x_3, x_4\}$, $\{y_1, x_3, y_4\}$, and $\{y_1, x_3, y_3\}$. Let $\Gamma_1, \Gamma_2, \dots, \Gamma_{n+1}$ be $\gamma(G)$ -sets containing y_1 , where $\{x_{n-1}, y_{n-1}\} \subseteq \Gamma_{n+1}$. Then, for $1 \leq i \leq n$, each Γ_i extends to $\Gamma'_i = \Gamma_i \cup \{x_{n+2}\}$ if $y_n \in \Gamma_i$ and $\Gamma'_i = \Gamma_i \cup \{y_{n+2}\}$ if $x_n \in \Gamma_i$, where each Γ'_i ($1 \leq i \leq n$) is a $\gamma(G')$ -set; $\Gamma_{n+1} = \{x_i, y_j \mid i \equiv 3, j \equiv 1 \pmod{4} \text{ and } 1 \leq i, j \leq n-2\} \cup \{x_{n-1}, y_{n-1}\}$ does not extend to a $\gamma(G')$ -set, but there exists a $\gamma(G')$ -set $\Gamma'_{n+1} = \{x_i, y_j \mid i \equiv 3, j \equiv 1 \pmod{4} \text{ and } 1 \leq i, j \leq n\} \cup \{x_{n+1}, y_{n+1}\}$ which does not come from any $\gamma(G)$ -set. Further, there exist two additional $\gamma(G')$ -sets which do not come from any $\gamma(G)$ -sets such as $\Gamma'_{n+2} = \{x_i, y_j \mid i \equiv 3, j \equiv 1 \pmod{4} \text{ and } 1 \leq i, j \leq n+1\} \cup \{x_{n+2}\}$ and $\Gamma'_{n+3} = \{x_i, y_j \mid i \equiv 3, j \equiv 1 \pmod{4} \text{ and } 1 \leq i, j \leq n+1\} \cup \{y_{n+2}\}$. So, noting that n is even, we have the following:

$$DV_{G'}^{2,2}(x_i) = \begin{cases} DV_G^{2,2}(x_i) & \text{if } i \equiv 0, 1, 2 \pmod{4} \text{ and } 1 \leq i \leq n-2 \\ DV_G^{2,2}(x_i) + 2 & \text{if } i \equiv 3 \pmod{4} \text{ and } 1 \leq i \leq n-2, \end{cases}$$

$$DV_{G'}^{2,2}(x_{n-1}) = \begin{cases} DV_G^{2,2}(x_{n-1}) + 2 & \text{if } n \equiv 0 \pmod{4} \\ DV_G^{2,2}(x_{n-1}) - 1 & \text{if } n \equiv 2 \pmod{4}, \end{cases}$$

$$DV_{G'}^{2,2}(x_{n+1}) = \begin{cases} 1 & \text{if } n \equiv 0 \pmod{4} \\ 3 & \text{if } n \equiv 2 \pmod{4}, \end{cases}$$

$$DV_{G'}^{2,2}(x_n) = DV_G^{2,2}(x_n), \text{ and } DV_{G'}^{2,2}(x_{n+2}) = \frac{n}{2} + 1.$$

Next, assume that (4) holds for G . Noting that $DV^2(v) = DV^{2,1}(v) + DV^{2,2}(v)$ and that n is even, by Subcase 2.1 and Subcase 2.2, we have

$$DV_{G'}^2(x_i) = \begin{cases} DV_G^2(x_i) & \text{if } i \equiv 0, 2 \pmod{4} \text{ and } 1 \leq i \leq n-2 \\ DV_G^2(x_i) + 2 & \text{if } i \equiv 1, 3 \pmod{4} \text{ and } 1 \leq i \leq n-2, \end{cases}$$

$$DV_{G'}^2(x_{n-1}) = DV_G^2(x_{n-1}) + 1, \quad DV_{G'}^2(x_n) = DV_G^2(x_n), \quad DV_{G'}^2(x_{n+1}) = 4, \text{ and } DV_{G'}^2(x_{n+2}) = n + 2, \text{ proving (4).}$$

Now, noting that $DV(v) = DV^1(v) + DV^2(v)$ for $v \in V(P_2 \square P_n)$, where $n \geq 4$ is even and $n \neq 6$, combine (3) and (4) to obtain (1), proving (ii).

(iii) By Theorem 3.4, $P_2 \square P_6$ has an additional γ -set $\{x_2, y_2, x_5, y_5\}$. This, together with (1), for $x_i, y_i \in V(P_2 \square P_6)$, we obtain

$$DV(x_i) = DV(y_i) = \begin{cases} 8 - i & \text{if } i \text{ is odd and } 1 \leq i \leq 3 \\ 5 & \text{if } i = 2 \text{ or } i = 5 \\ i + 1 & \text{if } i \text{ is even and } 4 \leq i \leq 6, \end{cases}$$

which equals the domination value in (2). \square

4 Total number of minimum dominating sets and domination value in $P_2 \square C_n$

For $n \geq 3$, consider $P_2 \square C_n$ as two copies of C_n with vertices labeled x_1, x_2, \dots, x_n and y_1, y_2, \dots, y_n with only the edges $x_i y_i$, for each i ($1 \leq i \leq n$), between two cycles (see Figure 4).




Figure 4: Labeling of vertices of $P_2 \square C_n$

We recall the following result.

Theorem 4.1. [7] For $n \geq 3$,

$$\gamma(P_2 \square C_n) = \begin{cases} \frac{n}{2} & \text{if } n \equiv 0 \pmod{4} \\ \lceil \frac{n+1}{2} \rceil & \text{if } n \not\equiv 0 \pmod{4}. \end{cases}$$

We introduce the following definition which will be used in the proof of Theorem 4.3.

Definition 4.2. Let G^1 and G^2 be disjoint copies of a graph G , and let D be a $\gamma(P_2 \square G)$ -set. Let $\langle D \cap V(G^1) \rangle = \cup_{i=1}^{m_1} \mathcal{H}_i^1$, a disjoint union of connected components such that $|V(\mathcal{H}_i^1)| \leq |V(\mathcal{H}_{i+1}^1)|$ for $1 \leq i \leq m_1 - 1$; similarly, we write $\langle D \cap V(G^2) \rangle = \cup_{i=1}^{m_2} \mathcal{H}_i^2$. Let $\alpha = \max(|V(\mathcal{H}_{m_1}^1)|, |V(\mathcal{H}_{m_2}^2)|)$; we will denote by \mathcal{H}_α any \mathcal{H}_i^j with $|V(\mathcal{H}_i^j)| = \alpha$, for $j = 1, 2$ ($1 \leq i \leq m_1$ or $1 \leq i \leq m_2$).

Example. The black vertices in Figure 5 form a $\gamma(P_2 \square C_{10})$ -set D , where $\langle D \rangle$ contains $2\mathcal{H}_2$.




Figure 5: $2\mathcal{H}_2 \subseteq \langle D \rangle$, where D is a $\gamma(P_2 \square C_{10})$ -set

Theorem 4.3. Let $n \geq 3$. For each $v \in V(P_2 \square C_n)$,

$$DV(v) = \begin{cases} 1 & \text{if } n \equiv 0 \pmod{4} \\ \frac{n+1}{2} & \text{if } n \equiv 1, 3 \pmod{4} \text{ and } n \neq 3 \\ (\lceil \frac{n+1}{2} \rceil)^2 & \text{if } n \equiv 2 \pmod{4} \text{ and } n \neq 6 \\ 3 & \text{if } n = 3 \\ 17 & \text{if } n = 6. \end{cases}$$

Proof. By Observation 2.2 and the vertex-transitivity, $DV(v) = DV(x_1)$ for each $v \in V(P_2 \square C_n)$. Let D be a $\gamma(P_2 \square C_n)$ -set containing x_1 , where $n \geq 3$; note that at least a vertex in $\{x_2, x_3, y_1, y_2, y_3\}$ belongs to D . Noting that each vertex dominates four vertices, we consider four cases.

Case 1. $n = 4k$, where $k \geq 1$: Since $\gamma(P_2 \square C_{4k}) = 2k$ and $|V(P_2 \square C_{4k})| = 8k$, each vertex is dominated by exactly one vertex (i.e., no vertex is doubly dominated). Thus there is a unique D containing x_1 , i.e., $D = \{x_i, y_j \mid i \equiv 1, j \equiv 3 \pmod{4}\}$, and hence $DV(x_1) = 1$.

Case 2. $n = 4k + 1$, where $k \geq 1$: Here $\gamma(P_2 \square C_{4k+1}) = 2k + 1$. We will show that no D contains both x_1 and a vertex in $\{y_1, y_2, x_3\}$. First, we note that no D contains both x_1 and y_1 : if $\{x_1, y_1\} \subseteq D$, then the part of $P_2 \square C_{4k+1}$ not dominated by $\{x_1, y_1\}$ is a $P_2 \square P_{4k-2}$, and $2k - 1$ vertices of $D - \{x_1, y_1\}$ must dominate $P_2 \square P_{4k-2}$. But $\gamma(P_2 \square P_{4k-2}) = 2k$ by Theorem 3.1, and we reach a contradiction. Second, we note that no D contains both x_1 and y_2 : if $\{x_1, y_2\} \subseteq D$, then the part of $P_2 \square C_{4k+1}$ not dominated by $\{x_1, y_2\}$ is the graph H in Figure 6, and $2k - 1$ vertices of $D - \{x_1, y_2\}$ must dominate H . If we let $S_0 = \{x_i, y_j \mid i \equiv 0, j \equiv$




Figure 6: $H \subset P_2 \square C_{4k+1}$

$2 \pmod{4}$ and $4 \leq i, j \leq 4k - 2\}$, then $|S_0| = 2(k - 1)$, S_0 dominates $8(k - 1)$ vertices, the part of H not dominated by S_0 is a P_4 , and one vertex of $D - (S_0 \cup \{x_1, y_2\})$ must dominate P_4 . But $\gamma(P_4) = 2$, and we reach a contradiction. (Similarly, no D contains both x_1 and y_{4k+1} .) Third, no D contains both x_1 and x_3 : if $\{x_1, x_3\} \subseteq D$, then a vertex in $N[y_2] = \{x_2, y_1, y_2, y_3\}$ must belong to D . Since $\{x_1, y_1\} \not\subseteq D$ (and thus $\{x_3, y_3\} \not\subseteq D$ by the vertex-transitivity) and $\{x_1, y_2\} \not\subseteq D$, $x_2 \in D$.

If $R_0 := \{x_1, x_2, x_3\} \subseteq D$, then the part of $P_2 \square C_{4k+1}$ not dominated by R_0 , say H_1 , must be dominated by $2k - 2$ vertices in $D - R_0$. Since $|V(P_2 \square C_{4k+1})| = 8k + 2$ and $|N[R_0]| = 8$, $2k - 2$ vertices in $D - R_0$ must dominate $8k - 6$ vertices. But each vertex in $P_2 \square C_{4k+1}$ dominates four vertices, and we reach a contradiction. (Similarly, no D contains both x_1 and x_{4k} .) So, we only need to consider D such that (i) $\{x_1, x_2\} \subseteq D$ (resp. $\{x_1, x_{4k+1}\} \subseteq D$) or (ii) no vertex in $N[x_1]$ is doubly dominated (i.e., $\{x_1, y_3\} \subseteq D$ and $\{x_1, y_{4k}\} \subseteq D$).

Subcase 2.1. $\{x_1, x_2\} \subseteq D$ (resp. $\{x_1, x_{4k+1}\} \subseteq D$): The part of $P_2 \square C_{4k+1}$ not dominated by $\{x_1, x_2\}$, say H_2 , must be dominated by $2k - 1$ vertices in $D - \{x_1, x_2\}$. Since $|V(P_2 \square C_{4k+1})| = 8k + 2$ and $|N[\{x_1, x_2\}]| = 6$, $2k - 1$ vertices in $D - \{x_1, x_2\}$ must dominate H_2 with $|V(H_2)| = 8k - 4$, and thus there exists at most one γ -set containing both x_1 and x_2 (resp. x_1 and x_{4k+1}). Noting that $\{x_1\} \cup \{x_i, y_j \mid i \equiv 2, j \equiv 0 \pmod{4}\}$ (resp. $\{x_i, y_j \mid i \equiv 1, j \equiv 3 \pmod{4}\}$) is a γ -set, there is a unique D containing both x_1 and x_2 (resp. x_1 and x_{4k+1}).

Subcase 2.2. No vertex in $N[x_1]$ is doubly dominated: Since $x_1 \notin V(\mathcal{H}_2)$, by Subcase 2.1, there are $2k - 1$ slots in which \mathcal{H}_2 can be placed.

By Subcase 2.1 and Subcase 2.2, we have $DV(x_1) = 2(1) + (2k - 1) = 2k + 1$.

Case 3. $n = 4k + 2$, where $k \geq 1$: Here $\gamma(P_2 \square C_{4k+2}) = 2k + 2$. We will show that no D contains a \mathcal{H}_α for $\alpha \geq 4$. If $R_1 := \{x_1, x_2, x_3, x_4\} \subseteq D$, then the part of $P_2 \square C_{4k+2}$ not dominated by R_1 , say F_1 , must be dominated by $2k - 2$ vertices in $D - R_1$. Since $|V(P_2 \square C_{4k+2})| = 8k + 4$ and $|N[R_1]| = 10$, $2k - 2$ vertices in $D - R_1$ must dominate F_1 with $|V(F_1)| = 8k - 6$. But each vertex in $P_2 \square C_{4k+2}$ dominates four vertices, and we reach a contradiction. We consider four subcases.

Subcase 3.1. $\mathcal{H}_3 \subseteq \langle D \rangle$: We denote by $DV^1(x_1)$ the number of such D 's containing x_1 . We note that the placement of \mathcal{H}_3 uniquely determines D : if $R_2 := \{x_1, x_2, x_3\} \subseteq D$, then the part of $P_2 \square C_{4k+2}$ not dominated by R_2 , say F_2 , must be dominated by $2k - 1$ vertices in $D - R_2$. Since $|V(P_2 \square C_{4k+2})| = 8k + 4$ and $|N[R_2]| = 8$, $2k - 1$ vertices in $D - R_2$ must dominate F_2 with $|V(F_2)| = 8k - 4$, and thus there exists at most one γ -set containing R_2 . Noting that $\{x_1, x_2\} \cup \{x_i, y_j \mid i \equiv 3, j \equiv 1 \pmod{4} \text{ and } 3 \leq i, j \leq 4k + 2\}$ is a γ -set, there is a unique D containing R_2 . If $x_1 \in V(\mathcal{H}_3)$, there are three such D 's, i.e., $\{x_1, x_2, x_3\} \subseteq D$, $\{x_{4k+2}, x_1, x_2\} \subseteq D$, and $\{x_{4k+1}, x_{4k+2}, x_1\} \subseteq D$. If $x_1 \notin V(\mathcal{H}_3)$, there are $2k - 1$ slots in which \mathcal{H}_3 can be placed. So, $DV^1(x_1) = 3 + (2k - 1) = 2k + 2$.

Subcase 3.2. $2\mathcal{H}_2 \subseteq \langle D \rangle$: We denote by $DV^2(x_1)$ the number of such D 's containing x_1 . Since each vertex in \mathcal{H}_2 is doubly dominated, four

vertices in $2\mathcal{H}_2$ are doubly dominated, and hence the placement of $2\mathcal{H}_2$ uniquely determines D . If $x_1 \in V(\mathcal{H}_2)$ (i.e., $\{x_1, x_2\} \subseteq D$ or $\{x_1, x_{4k+2}\} \subseteq D$), then there are $2k-1$ available slots to place the other \mathcal{H}_2 . If $x_1 \notin V(\mathcal{H}_2)$, then there are $\binom{2k-1}{2}$ available slots to place $2\mathcal{H}_2$'s. Thus, $DV^2(x_1) = 2(2k-1) + \binom{2k-1}{2} = (2k-1)(k+1)$.

Subcase 3.3. $\mathcal{H}_2 \subseteq \langle D \rangle$ and $2\mathcal{H}_2 \not\subseteq \langle D \rangle$: We will show that no such D exists. Without loss of generality, suppose that $\{x_1, x_2\} \subseteq D$. In order for y_3 to be dominated, a vertex in $N[y_3] = \{x_3, y_2, y_3, y_4\}$ must be in D . By the hypothesis, $\{x_1, x_2, x_3\} \not\subseteq D$. First, suppose that $R_3 := \{x_1, x_2, y_2\} \subseteq D$. Then the part of $P_2 \square C_{4k+2}$ not dominated by R_3 , say F_3 , must be dominated by $2k-1$ vertices in $D - R_3$. Since $|V(P_2 \square C_{4k+2})| = 8k+4$ and $|N[R_3]| = 7$, $2k-1$ vertices in $D - R_3$ must dominate F_3 with $|V(F_3)| = 8k-3$. But each vertex in $P_2 \square C_{4k+2}$ dominates four vertices, and we reach a contradiction. Second, suppose that $R_4 := \{x_1, x_2, y_3\} \subseteq D$. Then the part of $P_2 \square C_{4k+2}$ not dominated by R_4 , say F_4 , is a graph isomorphic to H in Figure 6, and $2k-1$ vertices of $D - R_4$ must dominate $F_4 \cong H$, which is a contradiction by Case 2. Third, suppose that $R_5 := \{x_1, x_2, y_4\} \subseteq D$. Then the part of $P_2 \square C_{4k+2}$ not dominated by R_5 , say F_5 , must be dominated by $2k-1$ vertices in $D - R_5$. Since $|V(P_2 \square C_{4k+2})| = 8k+4$ and $|N[R_5]| = 10$, $2k-1$ vertices in $D - R_5$ must dominate F_5 with $|V(F_5)| = 8k-6$, and thus there exist two vertices in $N[F_5]$ that are doubly dominated. When $k = 1$, one can easily see that $y_5 \in D$ (i.e., $2\mathcal{H}_2 \subseteq \langle D \rangle$) or $x_6 \in D$ (i.e., $\mathcal{H}_3 \subseteq \langle D \rangle$); both cases contradict to the assumption. So we consider for $k \geq 2$. Without loss of generality, we may assume that at least one vertex in $N[y_4] \cap N[F_5] = \{x_4, y_5\}$ is doubly dominated. In order for x_4 to be doubly dominated, $x_5 \in D$. If $\{x_1, x_2, y_4, x_5\} \subseteq D$, then the part of $P_2 \square C_{4k+2}$ not dominated by $\{x_1, x_2, y_4, x_5\}$ is the graph H' in Figure 7, and $2k-2$ vertices of $D - \{x_1, x_2, y_4, x_5\}$ must dominate H' . If we




Figure 7: $H' \subset P_2 \square C_{4k+2}$, where $k \geq 2$

let $S' = \{x_i, y_j \mid i \equiv 1, j \equiv 3 \pmod{4} \text{ and } 6 \leq i, j \leq 4k\}$, then $|S'| = 2k-3$, S' dominates $8k-12$ vertices, the part of H' not dominated by S' is a P_4 , and one vertex of $D - (S' \cup \{x_1, x_2, y_4, x_5\})$ must dominate P_4 . But $\gamma(P_4) = 2$ and we reach a contradiction. In order for y_5 to be doubly dominated, a vertex in $\{x_5, y_5, y_6\}$ must belong to D . Since $\{x_1, x_2, y_4, x_5\} \not\subseteq D$ and $\{x_1, x_2, y_4, y_5\} \not\subseteq D$, $y_6 \in D$. In this case, i.e.,

$\{x_1, x_2, y_4, y_6\} \subseteq D$, note that x_1 , x_2 , and y_5 are doubly dominated. In order for x_5 to be dominated, a vertex in $N[x_5] = \{x_4, x_5, x_6, y_5\}$ must be in D and each case results in at least two additional vertices to be doubly dominated, which is a contradiction. Thus, there is no $\gamma(P_2 \square C_{4k+2})$ -set containing exactly one \mathcal{H}_2 .

Subcase 3.4. $\mathcal{H}_2 \not\subseteq \langle D \rangle$: We denote by $DV^3(x_1)$ the number of such D 's containing x_1 . First, suppose that $\{x_s, y_s\} \subseteq D$ for some s ($1 \leq s \leq 4k+2$). If $\{x_1, y_1\} \subseteq D$, then the part of $P_2 \square C_{4k+2}$ not dominated by $\{x_1, y_1\}$ is $P_2 \square P_{4k-1}$, and $2k$ vertices of $D - \{x_1, y_1\}$ must dominate $P_2 \square P_{4k-1}$. By Theorem 3.4, there exist two such D 's for $k \neq 1$ (i.e., $n \neq 6$) and there exist three such D 's for $k = 1$ (i.e., $n = 6$). If $x_1 \in D$ and $\{y_1, y_2, y_{4k+2}\} \cap D = \emptyset$, then there are $2k$ available slots in which $\{x_s, y_s\} \subseteq D$ can be placed for some $s \neq 1$. Second, suppose that no two adjacent vertices belong to D . If we let $S_1 = \{x_i, y_j \mid i \equiv 1, j \equiv 3 \pmod{4} \text{ and } 1 \leq i, j \leq 4k\}$, then $|S_1| = 2k$ and the part of $P_2 \square C_{4k+2}$ not dominated by S_1 is a P_4 , so two vertices of $D - S_1$ must dominate P_4 . Since no two adjacent vertices belong to D , if $S_1 \subseteq D$, then $\{x_{4k}, y_{4k+1}\} \subseteq D$ or $\{x_{4k}, y_{4k+2}\} \subseteq D$ or $\{x_{4k+1}, y_{4k+2}\} \subseteq D$, thus there are two pairs of vertices (not necessarily disjoint) in D that are at distance two apart. The number of ways of selecting 2 out of $2k+2$ available slots is $\binom{2k+2}{2} = (k+1)(2k+1)$. Thus, $DV^3(x_1) = 2 + 2k + (k+1)(2k+1) = (k+1)(2k+3)$ if $k \neq 1$, and $DV^3(x_1) = 11$ if $k = 1$.

Now, noting that $DV(x_1) = DV^1(x_1) + DV^2(x_1) + DV^3(x_1)$, we have $DV(x_1) = (2k+2)^2$ if $k \neq 1$, and $DV(x_1) = 17$ if $k = 1$.

Case 4. $n = 4k + 3$, where $k \geq 0$: Here $\gamma(P_2 \square C_{4k+3}) = 2k + 2$. When $k = 0$, one can easily check that there are three γ -sets containing x_1 , i.e., $\{x_1, y_1\}$, $\{x_1, y_2\}$, and $\{x_1, y_3\}$. So $DV(x_1) = 3$ for $x_1 \in V(P_2 \square C_3)$. Next, we consider for $k \geq 1$. We will show that no D contains both x_1 and a vertex in $\{y_1, x_2, x_3\}$. First, note that no D contains both x_1 and y_1 : If $\{x_1, y_1\} \subseteq D$, then the part of $P_2 \square C_{4k+3}$ not dominated by $\{x_1, y_1\}$ is $P_2 \square P_{4k}$, and $2k$ vertices of $D - \{x_1, y_1\}$ must dominate $P_2 \square P_{4k}$. But $\gamma(P_2 \square P_{4k}) = 2k + 1$ by Theorem 3.1, and we reach a contradiction. Second, note that no D contains both x_1 and x_2 : if $\{x_1, x_2\} \subseteq D$, then the part of $P_2 \square C_{4k+3}$ not dominated by $\{x_1, x_2\}$, say H^* , must be dominated by $2k$ vertices. If we let $S^* = \{x_i, y_j \mid i \equiv 2, j \equiv 0 \pmod{4} \text{ and } 4 \leq i, j \leq 4k\}$, then $|S^*| = 2k - 1$ and the part of $P_2 \square C_{4k+3}$ not dominated by $S^* \cup \{x_1, x_2\}$ is a P_4 , and one vertex of $D - (S^* \cup \{x_1, x_2\})$ must dominate P_4 . But $\gamma(P_4) = 2$, and we reach a contradiction. (Similarly, no D contains both x_1 and x_{4k+3} .) Third, note that no D contains both x_1 and x_3 : if $\{x_1, x_3\} \subseteq D$, then a vertex in $N[y_2] = \{x_2, y_1, y_2, y_3\}$ must belong to D . Since $\{x_1, y_1\} \not\subseteq D$, $\{x_3, y_3\} \not\subseteq D$, and $\{x_1, x_2\} \not\subseteq D$,

we need to consider $\{x_1, x_3, y_2\} \subseteq D$: since $|V(P_2 \square C_{4k+3})| = 8k + 6$ and $|N[\{x_1, y_2, x_3\}]| = 8$, $2k - 1$ vertices of $D - \{x_1, x_3, y_2\}$ must dominate $8k - 2$ vertices, which is impossible since each vertex in $P_2 \square C_{4k+3}$ dominates four vertices. (Similarly, $\{x_1, x_{4k+2}\} \not\subseteq D$.) So, we only need to consider D such that (i) $\{x_1, y_2\} \subseteq D$ (resp. $\{x_1, y_{4k+3}\} \subseteq D$) or (ii) no vertex in $N[x_1]$ is doubly dominated. So suppose that $\{x_1, y_2\} \subseteq D$. Then the part of $P_2 \square C_{4k+3}$ that are not dominated by $\{x_1, y_2\}$, say H'' , must be dominated by $2k$ vertices. Since $|V(P_2 \square C_{4k+3})| = 8k + 6$ and $|N[\{x_1, y_2\}]| = 6$, $2k$ vertices of $D - \{x_1, y_2\}$ must dominate H'' with $|V(H'')| = 8k$, and thus there exists at most one such D . Since $\{x_1, y_2\} \cup \{x_i, y_j \mid i \equiv 0, j \equiv 2 \pmod{4} \text{ and } 3 \leq i, j \leq 4k + 3\}$ is a γ -set, if $\{x_1, y_2\} \subseteq D$, then there exists a unique such D . Similarly, there exists a unique D containing both x_1 and y_{4k+3} . If no vertex in $N[x_1]$ is doubly dominated (i.e., $\{x_1, y_3, y_{4k+2}\} \subseteq D$), then there are $2k$ slots in which a pair of vertices of D at distance two apart can be placed. Thus, $DV(x_1) = 2 + 2k$ if $k \geq 1$, and $DV(x_1) = 3$ if $k = 0$. \square

As an immediate consequence of Theorem 4.3, Observation 2.1, Observation 2.2, and the vertex-transitivity of $P_2 \square C_n$, we have the following.

Corollary 4.4. *For $n \geq 3$,*

$$\tau(P_2 \square C_n) \begin{cases} 4 & \text{if } n \equiv 0 \pmod{4} \\ 2n & \text{if } n \equiv 1, 3 \pmod{4} \text{ and } n \neq 3 \\ n(n+2) & \text{if } n \equiv 2 \pmod{4} \text{ and } n \neq 6 \\ 9 & \text{if } n = 3 \\ 51 & \text{if } n = 6. \end{cases}$$

5 Open Problems

We end this paper with some open problems. One could ask the following questions.

1. In our terminology, Mynhardt [16] characterized vertices v in a tree T such that $DV(v) = \tau(T)$ or $DV(v) = 0$. Can we describe vertices satisfying $DV(v) = k$ for $k \neq 0, \tau(T)$?
2. For $e \in E(G)$, can we find the bounds of $\tau(G - e)$ in terms of $\tau(G)$? And, for $v \in V(G - e)$, how does $DV_{G-e}(v)$ change in terms of $DV_G(v)$?
3. For $w \in V(G)$, can we find the bounds of $\tau(G - w)$ in terms of $\tau(G)$? And, for $v \in V(G - w)$, how does $DV_{G-w}(v)$ change in terms of $DV_G(v)$?
4. For a given graph G , can we characterize subgraphs $H \subseteq G$ satisfying $DV_H(v) = DV_G(v)$ for each vertex $v \in V(H)$?

In parallel with the idea of $\tau(G)$, the anonymous referee suggested the following questions.

5. Can we compute the *number* of *ir*-sets (maximal irredundant sets of minimum cardinality), γ -sets (minimum dominating sets), γ_t -sets (minimum total dominating sets), *i*-sets (minimum independent dominating sets), β_0 -sets (maximum independent sets), Γ -sets (minimal dominating sets of maximum cardinality), *IR*-sets (maximum irredundant sets) in a graph G ?

Acknowledgement. The author thanks Cong X. Kang for suggesting the notion of domination value, as well as his helpful comments and suggestions. The author also thanks the referee for bringing to her attention the reference [18], and other helpful comments and suggestions.

References

- [1] C. Berge, *Graphs and Hypergraphs* (North-Holland, Amsterdam, 1973).
- [2] C. Berge, *Theory of Graphs and its Applications* (Methuen, London, 1962).
- [3] T. Y. Chang and W. E. Clark, The domination numbers of the $5 \times n$ and $6 \times n$ grid graphs. *J. Graph Theory* **17** No. 1 (1993), 81-107.
- [4] G. Chartrand and P. Zhang, *Introduction to Graph Theory* (McGraw-Hill, Kalamazoo, MI, 2004).
- [5] E. J. Cockayne, R. M. Dawes, and S. T. Hedetniemi, Total domination in Graphs. *Networks* **10** (1980), 211-219.
- [6] E. J. Cockayne and S. T. Hedetniemi, Towards a Theory of Domination in Graphs. *Networks* **7** (1977), 247-261.
- [7] L. Eroh, R. Gera, C. X. Kang, C. E. Larson, and E. Yi, Domination in Functigraphs. *Discuss. Math. Graph Theory*, to appear.
- [8] E. O. Hare, Algorithms for grid and grid-like graphs. Ph. D. Dissertation (Clemson University, 1989).
- [9] T. W. Haynes, S. T. Hedetniemi, and P. J. Slater, *Fundamentals of Domination in Graphs* (Marcel Dekker, New York, 1998).
- [10] T. W. Haynes, S. T. Hedetniemi, and P. J. Slater, editors, *Domination in Graphs: Advanced Topics* (Marcel Dekker, New York, 1998).

- [11] S. T. Hedetniemi and R.C. Laskar, Introduction, *Discrete Math.* **86** (1990), 3-9.
- [12] M. A. Henning, A survey of selected recent results on total domination in graphs. *Discrete Math.* **309**, Issue 1 (2009), 32-63.
- [13] M. S. Jacobson and L. F. Kinch, On the domination number of products of graphs I. *Ars Combin.* **18** (1983), 33-44.
- [14] C. X. Kang, Total Domination Value in Graphs. *Util. Math.*, to appear.
- [15] S. Klavžar and N. Seifter, Dominating Cartesian products of cycles. *Discrete Appl. Math.* **59** (1995), 129-136.
- [16] C. M. Mynhardt, Vertices Contained in Every Minimum Dominating Set of a Tree. *J. of Graph Theory* **31** (1999), 163-177.
- [17] O. Ore, *Theory of Graphs* (Amer. Math. Soc. Colloq. Publ., **38**, Providence, 1962).
- [18] P. J. Slater, The Hedetniemi number of a graph. *Congr. Numer.* **139** (1999), 65-75.
- [19] E. Yi, Domination Value in Graphs. *submitted*.